

Volume 19 No. 1

MGH CARES ABOUT PAIN RELIEF

January 2020

PAIN RELIEF CONNECTION THE PAIN INFORMATION NEWSLETTER

Provided by MGH Cares About Pain Relief, a program of MGH Patient Care Services

Inside this issue:		Complementary Integrative Health	Page 2
In the News	Page 1	Pain Resources on the Web	Page 2
Journal Watch	•	Pain Education Opportunities	Page 2
o Nows	U	MGH Pain Calendar & Resources	Page 2

In the News

- CDC data shows Massachusetts with the 4th lowest opioid prescribing rate has the 9th highest rate of overdoses in America.
- FDA warns about <u>breathing problems with gabapentinoids</u> for neuropathic pain in those with respiratory comorbidities or CNS depressants.
- Dr. Ronald Melzack pioneer in the field of pain (Gate & Neuromatrix) theory, education and research passed away December 22nd 2019.
- Massachusetts releases its Electronic Prescribing and Dispensing Manual, a new requirement for controlled substances.
- Duexis (ibuprofen/famotidine) & Vimovo (Naproxen/Esomeprazole) under fire as examples of price gouging payers.
- Injectable ibuprofen, now in infusion bags ready to use for adults and children over 6 months old.
- World Health Organization has revised the WHO Guideline on the Management of Chronic Pain in Children
- Opioids were prescribed to 41% fewer adults with moderate or severe pain between 2014 and 2016.

Journal Watch [MGHers can obtain articles through the Treadwell home page]

- National Academies of Sciences, Engineering, and Medicine. (2019) Framing Opioid Prescribing Guidelines for Acute Pain: Developing the Evidence. Washington, DC: The National Academies Press. See <u>full document</u> or <u>highlights</u> developing a model to reduce overprescribing.
- Oderda GM, Senagore AJ, Morland K, et al. Opioid-related respiratory and gastrointestinal adverse events in patients with acute postoperative pain: prevalence, predictors, and burden. *J Pain Palliat Care Pharmacother*. 2019; 33(3-4):82-97. Prevalence of opioid-induced respiratory (3-17%) & GI (42-72%) adverse events is high; affecting outcomes. At risk patients, doses & procedures are detailed.
- Bout-Tabaku S, Gupta R, Jenkins TM. Musculoskeletal pain, physical function, and quality of life after bariatric surgery. *Pediatrics*. 2019;144(6):e20191399. Morbidly obese teens had 40% less pain within a year following bariatric surgery.
- Saragiotto BT, et al. Paracetamol for pain in adults. BMJ. 2019;367:16693. Safety & efficacy of acetaminophen for chronic pain isn't known.
- Kube T, Rief W, Vivell MB, et al. Deceptive and Nondeceptive Placebos to Reduce Pain: An Experimental Study in Healthy Individuals. *Clin J Pain*. 2020;36:68-79. Open label placebo doesn't cut pain intensity or unpleasantness but may improve its tolerability in healthy people.
- Igler E, Lang A, Balistreri K, et al. <u>Parents reliably identify pain dismissal</u> by pediatric providers. *Clin J Pain*. 2020 Feb;36(2):80-87. Parents listening provider interaction vignettes with a pediatric chronic pain patient rated the dismissive scenario more negatively. Gender played a role, but when asked what they would say to the dismissive provider, parents were likely to defend their child & ask for a second opinion.
- Tarn DM, Barrientos M, et al. Prevalence and knowledge of potential interactions between over-the-counter products and apixaban. *J Am Geriatr Soc.* 2020 Jan;68(1):155-162. Apixaban has bleeding risk from interactions with certain diet supplements or NSAIDs used for pain.

MGH Cares About Pain Relief
Massachusetts General Hospital

Previous Newsletters

Journal Watch (continued) [MGHers can obtain articles through the Treadwell home page]

- Beal SJ, Kashikar-Zuck S, King C, et al. Heightened risk for pain in young adult women with a history of childhood maltreatment: a
 prospective longitudinal study. Pain. 2020 Jan;161(1):156-65. Women who endured childhood maltreatment reported a greater number of
 pain locations, more intensity & more frequent episodes of pain than matched women who were not maltreated as a child or adolescent.
- Chua KP, Brummett CM, Conti RM, et al. Association of opioid prescribing patterns with <u>prescription opioid overdose in adolescents and young adults</u>. *JAMA Pediatr*. 2019 Dec 16. Among 2.7 million youth prescribed opioids, 0.01% overdosed; with higher risk factors identified.
- Stearns LM, Abd-Elsayed A, Perruchoud C, et al. Intrathecal drug delivery systems for cancer pain: an analysis of a prospective, multicenter product surveillance registry. Anesth Analg. 2020 Feb;130(2):289-297. A study of >1,400 cancer pain patients showed at a year intrathecal pumps significantly improve pain, mobility, functioning & mood with 10% discontinuing, having complication or device removal.
- Voepel-Lewis T, Farley FA, Grant J, et al. Behavioral intervention and <u>disposal of leftover opioids</u>: a randomized trial. *Pediatrics*. 2020
 Jan;145(1). Parental education & providing a disposal method increased the likelihood that a child's leftover opioids were properly disposed.
- Sodhi J, Karmarkar A, Raji M, et al. Pain as a predictor of frailty over time among older Mexican Americans. *Pain.* 2020 Jan;161(1):109-13. Pain increases the risk of frailty by 70% in older Mexican Americans with age, gender, depressed mood & hip fractures also risk factors.

Pain Resources on the Web:

- Patients perceive more empathy & experience more pain relief when providers use therapies they believe will work.
- Use the <u>FDA's infographic on drug disposal methods</u> to educate patients and clarify confusion over mixed messages.
- Understanding pediatric cancer pain through qualitative research of the lived experience. Cancer changes the meaning of pain.
- There is a need to <u>understand & talk about the biopsychosocial nature of chronic pain</u> in a nonjudgmental & nonpejorative way.

Complementary Integrative Health

- Despite many studies show benefits of aromatherapy, massage & reflexology for palliation; research methods must be strengthened.
- Systematic review & meta-analysis shows <u>acupuncture and /or acupressure cuts pain & medication use for cancer patients.</u>
- Meditation, hypnosis, suggestion & cognitive behavioral therapy reduce pain & opioid use among best researched mind-body therapies.
- An anti-inflammatory diet is likely to reduce the pain hypersensitivity associated with pro-inflammatory diets in patients with fibromyalgia.

Pain-Related Education Opportunities

- Wed-Sun Feb 26 Mar 1st American Academy of Pain Medicine meeting covers a range of pain & treatment types in National Harbor, MD
- Mon. Mar 9th 10:30 11:30am Amygdala Pain Mechanisms Role of the CRF System: an NIH pain seminar
- Tue Mar 24th Management of Chronic Pain in an Opioid-Phobic Society (Drug & nondrug approaches) St. Anslem. Manchester NH

MGH Pain Calendar

- Tools & Techniques for Effective Pain Management Tue, April 28th FND 325. 4.5 CE with 1 pharmacology credit Click here to register.
- Palliative Care Grand Rounds are held Wednesdays from 8:00 AM 9:00 AM in the Ether Dome Email: Gail Chin for more information.
- MGH Inter-professional Pain Rounds Jackson 412 First and third Wednesday of month from 12–1pm. Email <u>Tina Toland</u> for more info.

MGH Pain Resources

Excellence Every Day Pain Portal Page: http://www.mghpcs.org/EED/Pain/default.shtml

The MGH Center for Translational Pain Research: https://www.massgeneral.org/TranslationalPainResearch/

MGH Palliative Care: http://www.massgeneral.org/palliativecare

MGH Formulary (includes patient teaching handouts in 16 languages): http://www.crlonline.com/crlsql/servlet/crlonline

MGH Cares About Pain Relief
Massachusetts General Hospital

Previous Newsletters

To be added to or removed from the Pain Relief Connection mailing list, or other inquiries, send an email to Paul Amstern